


# Lecture 7: Limits in several variables (Sect 14.2) (22)


Last time:  $E: \mathbb{R} \rightarrow \mathbb{R}$

Say  $\lim_{h \rightarrow 0} E(h) = 0$  if given  $\epsilon > 0$  can always find a  $\delta > 0$  so that when  $0 < |h| < \delta$  then  $|E(h)| < \epsilon$



[Foreshadow story of the Sleipner A sinking.]

Now suppose  $E: \mathbb{R}^2 \rightarrow \mathbb{R}$ . We say  $\lim_{\vec{h} \rightarrow 0} E(\vec{h}) = 0$  if given  $\epsilon > 0$  can always find  $\delta > 0$  so that when  $0 < |\vec{h}| < \delta$  then  $|E(\vec{h})| < \epsilon$ .




Ex:

$$\lim_{(x,y) \rightarrow (0,0)} x + 3y = 0$$

Reason: Let  $\epsilon > 0$  be given. Take  $\delta = \epsilon/4$ .

If  $h = (x, y)$  and  $|h| < \delta$ , then  $|x| < \delta$  and  $|y| < \delta$  as shown:



Then

$$|x+3y| \leq |x| + |3y| < \delta + 3\delta = 4\delta = \varepsilon.$$

More generally, if  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  then

$$\lim_{\vec{x} \rightarrow \vec{a}} f(\vec{x}) = c \text{ if } f(\vec{a} + \vec{h}) = c + E(\vec{h})$$

where  $\lim_{\vec{h} \rightarrow \vec{0}} E(\vec{h}) = 0.$

Same notion of limit works for  $f: \mathbb{R}^3 \rightarrow \mathbb{R}$   
or  $\mathbb{R}^n \rightarrow \mathbb{R}$  or even  $\mathbb{R}^n \rightarrow \mathbb{R}^m.$

A more complicated example:

$$\text{Take } f(x,y) = \frac{2xy}{x^2+y^2}$$

not defined  
at  $(0,0).$

What is

$$\lim_{(x,y) \rightarrow \vec{0}} f(x,y) = ?$$

[First, try our usual trick: reduce the dimension]

Along the x-axis:

$$f(x, 0) = \frac{2x \cdot 0}{x^2 + 0^2} = 0$$

which suggests  $\lim = 0$ . But along the line  $y = x$

$$f(x, x) = \frac{2x \cdot x}{x^2 + x^2} = 1 \quad (\text{for } x \neq 0)$$

Thus the limit does not exist.


Can actually work out the graph of  $f$ .


Along the line  $y = cx$  have

$$f(x, cx) = \frac{2x(cx)}{x^2 + (cx)^2} = \frac{2cx^2}{(1+c^2)x^2} = \frac{2c}{1+c^2}$$

E.g.  $f = -1$  along  $y = -x$

The full picture is:





Even odder:

$$f(x, y) = \frac{xy^2}{x^2 + y^4}$$


What is limit  
as  $(x, y) \rightarrow \vec{0}$ ?

Along the line  $y = cx$  we have

$$f(x, cx) = \frac{x(cx)^2}{x^2 + (cx)^4} = \frac{cx^3}{x^2(1+c^4x^2)} = \frac{cx}{1+c^4x^2}$$

which  $\rightarrow 0$  as  $x \rightarrow 0$

But: Along  $x = y^2$   
we have



$$f(y^2, y) = \frac{y^2 \cdot y^2}{(y^2)^2 + y^4} = \frac{y^4}{2y^4} = \frac{1}{2}.$$


So again, the limit does not exist!

Rules for limits:

$$f, g: \mathbb{R}^2 \rightarrow \mathbb{R}$$

a)  $\lim_{\vec{x} \rightarrow \vec{a}} (f(\vec{x}) + g(\vec{x}))$   
 $= \lim_{\vec{x} \rightarrow \vec{a}} f(\vec{x}) + \lim_{\vec{x} \rightarrow \vec{a}} g(\vec{x})$

b)  $\lim_{\vec{x} \rightarrow \vec{a}} f(\vec{x})g(\vec{x})$   
 $= \left( \lim_{\vec{x} \rightarrow \vec{a}} f(\vec{x}) \right) \left( \lim_{\vec{x} \rightarrow \vec{a}} g(\vec{x}) \right)$



In both cases, this is provided the RHS all makes sense.